

Institute of Natural and Life Sciences

Department of Biotechnology

Module: Biophysics

Series 4: Diffusion phenomena**Exercise: 1**

A porous membrane with a total pore surface area $S=0.05\text{m}^2$ separates two compartments containing sucrose at concentrations of 0.5 and 0.2 mol/L, respectively. These concentrations are maintained constant during the diffusion of sucrose molecules through the membrane. It is assumed that a steady-state regime is established.

- What is the value of the flow rate?

Given: diffusion coefficient of sucrose $D=8\times 10^{-10}\text{ m}^2/\text{s}$, membrane thickness $e=10\text{ }\mu\text{m}$.

Exercise: 2

Let there be a porous membrane with a thickness e and a surface area $S=50\text{cm}^2$ separating two compartments. At time $t=0$, 2 liters of pure water are introduced into the first compartment and 2 liters of an aqueous solution with a solute concentration of 1 mole/L are introduced into the second compartment. If after 30 seconds the concentration in the first compartment is $10^{-6}\text{mole}/\text{cm}^3$

- determine the thickness e of the membrane, assuming that the concentration gradient remains linear across the thickness e . Given: $D=5.344\times 10^{-5}\text{cm}^2/\text{s}$

Exercise: 3

The diffusion coefficient of insulin in aqueous solution at 25°C is equal to $8.2\times 10^{-11}\text{m}^2\text{s}^{-1}$.

1. Calculate the radius of this molecule, assumed to be spherical.
2. Deduce the molar mass of insulin from this result.
3. What would be the diffusion coefficient of insulin at 0°C ?
4. What would be the diffusion coefficient of urea in aqueous solution at 0°C ?

The given data is: The density of insulin: $1300\text{ kg}/\text{m}^3$, $\eta_{\text{H}_2\text{O}}$: $1\text{ mPa}\cdot\text{s}$, $K=1.38\times 10^{-23}\text{J}/\text{K}$,
 M_{urea} : $60\text{ g}/\text{mol}$

Exercise:4

A reservoir is divided into two compartments by a porous membrane with a surface area of 3 cm^2 and a thickness of 0.1 mm . In one of the compartments, an aqueous solution of $2\text{ mmol}/\text{L}$ is placed, and in the other, pure water. The initial molecular diffusion flux of the solute is $4.2\times 10^{-12}\text{mol}/\text{s}$.

1. Calculate the permeability coefficient P of the membrane for the molecule.
2. Deduce the molecular diffusion coefficient from this.