


CHAPTER 3. Respiration

I. Respiration in invertebrates

Respiration in invertebrates refers to the process of **gas exchange**, whereby oxygen is obtained from the environment and carbon dioxide is eliminated from the body. The mechanisms of respiration in these organisms are highly diverse and strongly influenced by **body size, level of activity, and environmental habitat** (aquatic vs. terrestrial). In general, invertebrates exhibit four principal strategies of respiration: **cutaneous exchange, gills, tracheal systems, and book lungs**.

2.1. Cutaneous Respiration


Cutaneous respiration involves the **direct diffusion of gases across the body surface**. This mechanism is effective only in organisms with a **thin body wall** and a **high surface-area-to-volume ratio**, allowing oxygen to diffuse efficiently into underlying tissues. It is common in **annelids (earthworms)** and **small flatworms (Platyhelminthes)**, which lack specialized respiratory organs. Because diffusion is a slow process, cutaneous respiration is generally associated with **small-sized organisms** or those living in **moist environments**, where the skin remains permeable to gases.



2.2. Gills (Branchiae)


In aquatic invertebrates, the most common respiratory structures are **gills (branchiae)**. These are thin, vascularized extensions of the body surface or appendages, adapted for gas exchange with water. Gills are present in many groups, including **mollusks, crustaceans, and some annelids**. The extensive surface area of gills, combined with their rich vascularization, facilitates efficient uptake of dissolved oxygen and elimination of carbon dioxide. In addition,

in many species, **cilia or specialized appendages** generate water currents to ensure continuous renewal of the water surrounding the gill surface, thereby maximizing respiratory efficiency.



2.3. Tracheal System


A unique adaptation found in terrestrial invertebrates, particularly **insects and some arachnids**, is the **tracheal system**. This system consists of a branching network of fine air-filled tubes (tracheae) that penetrate the body and deliver oxygen **directly to the tissues and cells**, bypassing the need for a circulatory transport of gases. The tracheae open to the exterior through small valves called **spiracles**, which can regulate air flow and water loss. This mode of respiration is highly efficient for small terrestrial organisms with high metabolic demands, such as insects, allowing them to sustain intense activity (e.g., flight).



2.4. Book Lungs

Certain arachnids, such as **spiders and scorpions**, possess specialized respiratory organs known as **book lungs**. These consist of internal chambers containing numerous thin, leaf-like lamellae stacked in parallel, resembling the pages of a book. Air enters the chamber through a small opening on the body surface and diffuses across the lamellae into the blood or

hemolymph. The lamellar structure provides a **large surface area for gas exchange**, making book lungs particularly suited for terrestrial life in environments with relatively low humidity.



2.5. Environmental Adaptations

The diversity of respiratory structures among invertebrates illustrates the importance of **environmental adaptation**.

- **Aquatic invertebrates** predominantly rely on **gills** or **cutaneous exchange**, optimized for extracting oxygen dissolved in water.
- **Terrestrial invertebrates**, in contrast, have evolved specialized systems such as **tracheae** or **book lungs**, enabling them to cope with air breathing and the risk of desiccation.


Thus, the type of respiratory mechanism employed by a given invertebrate group reflects a balance between **morphological constraints, ecological pressures, and metabolic requirements**.

II. Respiration in vertebrates

Respiration in vertebrates is the process of taking in oxygen (O_2) and removing carbon dioxide (CO_2). The organs and breathing methods change according to whether the animal lives in water or on land, but all respiratory surfaces must be **large, thin, moist, and rich in blood vessels** to allow gases to diffuse easily.


2.1. Gills

Fishes breathe mainly with **gills**. Gills are thin filaments covered with lamellae that provide a very large surface area for gas exchange. Water enters the mouth, flows over the gills, and exits through the gill openings or operculum. The fish pumps water by opening and closing its mouth. Blood inside the gill filaments flows in the **opposite direction to the water (countercurrent flow)**, which keeps a strong oxygen gradient and allows maximum oxygen uptake.




2.2. Skin and Lungs

Amphibians (like frogs) use both their **skin** and **lungs** for breathing. The skin is moist and thin, allowing oxygen to pass directly into blood vessels, especially when the animal is underwater. For lung breathing, amphibians use **positive-pressure breathing**. They lower the floor of the mouth to draw air in through the nostrils, then raise the floor to push air into the lungs. Air is forced in rather than sucked in.




2.3. Lungs

Reptiles rely mainly on **lungs**, which are more folded and efficient than amphibian lungs. They use **negative-pressure breathing**. Muscles between the ribs expand the chest cavity, lowering the pressure inside the lungs so air is drawn in. When the muscles relax, the chest cavity becomes smaller and air is pushed out.




2.4. Lungs with air sacs

Birds have the **most efficient respiratory system** of all vertebrates. Their rigid lungs are connected to several **air sacs** that act as bellows. Breathing occurs in **two cycles**. During the first inhalation, air moves into the posterior air sacs. During the first exhalation, fresh air passes through the lungs where gas exchange occurs. On the second inhalation, used air moves into the anterior air sacs. On the second exhalation, this air leaves the body. This system keeps **fresh air moving through the lungs during both inhalation and exhalation**, providing a constant supply of oxygen.



2.5. Lungs with Alveoli

Mammals breathe using highly branched **lungs** that end in millions of tiny **alveoli**, which greatly increase the surface area for gas exchange. Mammals use **negative-pressure breathing** controlled by the **diaphragm**. When the diaphragm contracts and moves downward and the rib cage expands, the chest cavity volume increases and pressure inside decreases, pulling air into the lungs. When the diaphragm relaxes, the chest volume decreases and air is pushed out.



Summary

Respiration in animals ensures the exchange of oxygen and carbon dioxide necessary for life. **Invertebrates** show diverse methods adapted to their environment: **cutaneous respiration** through the skin, **gills** in aquatic forms, **tracheal systems** in insects, and **book lungs** in arachnids. These mechanisms reflect adaptations to body structure and habitat. **Vertebrates** possess specialized organs that evolved for greater efficiency—**fish** use gills with countercurrent flow, **amphibians** breathe through skin and simple lungs, **reptiles** use more developed lungs, **birds** have lungs with air sacs for continuous airflow, and **mammals** possess lungs with millions of alveoli for maximal gas exchange. Overall, respiratory systems evolve from simple diffusion to highly efficient lungs to meet increasing metabolic demands.